



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Self-Interaction Free and Analytic Treatment of the Coulomb Energy in Kohn-Sham Density Functional Theory

M. Daene, A. Gonis, D. Nicholson, G. M. Stocks

February 27, 2012

APS March Meeting
Boston, MA, United States
February 27, 2012 through March 2, 2012

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

Self-Interaction Free and Analytic Treatment of the Coulomb Energy in Kohn-Sham Density Functional Theory

M. Daene^{1,2}, A. Gonis¹, D.M. Nicholson², G.M. Stocks²

¹) Lawrence Livermore National Laboratory, ²) Oak Ridge National Laboratory



Research at ORNL sponsored by the U.S. Department of Energy, Office of Basic Energy Sciences, Materials Sciences and Engineering Division, "Center for Defect Physics," an Energy Frontier Research Center.

LLNL-CONF-533132

This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under contract DE-AC52-07NA27344. Lawrence Livermore National Security, LLC



Overview

- The use of the classical Hartree term in implementations of KS-DFT leads to self-interaction effects.
- A great number of functionals has been developed over the years with the purpose of removing the effects of self-interaction from the theory.
- We have now solved uniquely, analytically and in closed form the self-interaction problem in Kohn-Sham Density Functional Theory.
- We will describe how this is done, and show results of calculations for realistic atomic systems and will compare with competing methodologies (OEP, exact exchange).
- **The underlying formalism is that of Density Function Theory(DFT), in which the independent variable is the density $n(r)$.**
- **Functional derivatives taken only with respect to the density**

Kohn-Sham Density Functional Theory

real system

$$\hat{H} = V_{ext} + \hat{T} + \hat{U}$$

$$\begin{aligned} E[n] &= \langle \Psi | \hat{H} | \Psi \rangle \\ &= \int V_{ext}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} + \langle \Psi | \hat{T} + \hat{U} | \Psi \rangle \end{aligned}$$

Hohenberg and Kohn have shown that

$$\left. \frac{\delta E[n]}{\delta n(\mathbf{r})} \right|_{n=n_{GS}} = 0$$

$$\left. \frac{\delta \langle \Psi | \hat{T} + \hat{U} | \Psi \rangle}{\delta n(\mathbf{r})} \right|_{n=n_{GS}, \Psi=\Psi_{GS}} = -V_{ext}$$

fictitious non-interacting system with the same density

$$\hat{H}_s = V_s + \hat{T}$$

$$\begin{aligned} E_s[n] &= \langle \Phi | V_s + \hat{T} | \Phi \rangle \\ &= \int V_s(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} + \langle \Phi | \hat{T} | \Phi \rangle \end{aligned}$$

$$\left. \frac{\delta E_s[n]}{\delta n(\mathbf{r})} \right|_{n=n_{GS}} = 0$$

$$\left. \frac{\delta \langle \Phi | \hat{T} | \Phi \rangle}{\delta n(\mathbf{r})} \right|_{n=n_{GS}, \Phi=\Phi_{GS}} = -V_s$$

$$\langle \Psi_{GS} | \hat{T} + \hat{U} | \Psi_{GS} \rangle \leq \langle \Phi_{GS} | \hat{T} + \hat{U} | \Phi_{GS} \rangle$$

Kohn-Sham Density Functional Theory

$$V_s(\mathbf{r}) = V_{ext}(\mathbf{r}) + \frac{\delta}{\delta n(\mathbf{r})} \langle \Phi_{GS} | \hat{U} | \Phi_{GS} \rangle + \frac{\delta E_c[n]}{\delta n(\mathbf{r})}$$

Kohn-Sham equations $[\hat{T} + V_s] f_i = \epsilon_i f_i$

Φ is a Slater Determinant build of f_i

pair density

$$U = \frac{1}{2} \int \int \frac{n^{(2)}(\mathbf{r}, \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r} d\mathbf{r}' = \frac{1}{2} \int \int \frac{n(\mathbf{r}) n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r} d\mathbf{r}' + \frac{1}{2} \int \int \frac{J(\mathbf{r}, \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r} d\mathbf{r}'$$

exchange term: $J(\mathbf{r}, \mathbf{r}') = - \sum_{ij} f_i^*(\mathbf{r}) f_j^*(\mathbf{r}') f_j(\mathbf{r}) f_i(\mathbf{r}') \delta_{\sigma_i, \sigma_j}$

The exchange term depends only implicitly on the density!

New Method of Calculating the Functional Derivative

- expand orbitals in orthonormal and complete basis
- basis written explicitly in term of the density
- differentiate the expansion (functional differentiation)
- get the potential

Equidensity Basis

new basis as functional of n : $\phi_{\mathbf{k}}(\mathbf{r}, [n]) = \sqrt{\frac{n(\mathbf{r})}{N}} \exp\{i \mathbf{k} \cdot \mathbf{R}(\mathbf{r}, [n])\}$

John E. Harriman, Phys. Rev. A 24, 680 (1981)

Gil Zumbach and Klaus Maschke, Phys. Rev. A 2, 544 (1983)

with

$$\mathbf{k} = (k_x, k_y, k_z)$$

$$R_1(\mathbf{r}) = R_1(x, y, z) = 2\pi \frac{\int_{-\infty}^x dx' n(x', y, z)}{\int_{-\infty}^{\infty} dx' n(x', y, z)}$$

$$R_2(\mathbf{r}) = R_2(y, z) = 2\pi \frac{\int_{-\infty}^y dy' \int_{-\infty}^{\infty} dx' n(x', y', z)}{\int_{-\infty}^{\infty} dy' \int_{-\infty}^{\infty} dx' n(x', y', z)}$$

$$R_3(\mathbf{r}) = R_3(z) = \frac{2\pi}{N} \int_{-\infty}^z dz' \int_{-\infty}^{\infty} dy' \int_{-\infty}^{\infty} dx' n(x', y', z')$$







The equidensity basis is complete and orthonormal (in \mathbf{r}) for any density !!!

a representation of orbitals in this basis is possible


$$f_j(\mathbf{r}) = \sum_{\mathbf{k}} a_{\mathbf{k}}^j \phi_{\mathbf{k}}(\mathbf{r}, [n])$$

$$\frac{\delta f_i(\mathbf{r})}{\delta n(\mathbf{r}')} = \frac{\delta \mathbf{R}[n(\mathbf{r})]}{\delta n(\mathbf{r}')} \cdot \sum_{\mathbf{k}} a_{\mathbf{k}}^i i \mathbf{k} \phi_{\mathbf{k}}[n(\mathbf{r})] + \frac{\delta(\mathbf{r} - \mathbf{r}')}{2n(\mathbf{r})} f_i(\mathbf{r})$$







Exchange Potential: He and Be




Exchange Potential: Ne and Ar



Exchange Potential: Li and B



Exchange Potential: He to Kr




Energies – Atom Series

Z	Symbol	Name	Exp	x-SIF	x-OEP	HF
2	He	Helium	-2.904	-2.862	-2.862	-2.862
3	Li	Lithium	-7.47806	-7.432	-7.433	-7.433
4	Be	Beryllium	-14.66736	-14.571	-14.572	-14.573
5	B	Boron	-24.65391	-24.527	-24.528	-24.529
6	C	Carbon	-37.8450	-37.687	-37.689	-37.690
7	N	Nitrogen	-54.5892	-54.401	-54.403	-54.405
8	O	Oxygen	-75.0673	-74.809	-74.812	-74.814
9	F	Fluorine	-99.7339	-99.406	-99.409	-99.411
10	Ne	Neon	-128.9376	-128.542	-128.545	-128.547
11	Na	Sodium	-162.2546	-161.852	-161.857	-161.859
12	Mg	Magnesium	-200.053	-199.606	-199.612	-199.615
13	Al	Aluminum	-242.346	-241.868	-241.873	-241.877
14	Si	Silicon	-289.359	-288.845	-288.851	-288.854
15	P	Phosphorus	-341.259	-340.709	-340.715	-340.719
16	S	Sulfur	-398.110	-397.495	-397.502	-397.506
17	Cl	Chlorine	-460.148	-459.470	-459.478	-459.483
18	Ar	Argon	-527.540	-526.804	-526.812	-526.817

energies in H

we find: $E_g \leq E^{HF} \leq E^{OEP} \leq E^{SIF}$

Energies – Atom Series



Comparison to OEP

DFT condition for the ground state:

$$\frac{\delta E}{\delta n} = 0$$


OEP condition for the ground state:

$$0 = \frac{\delta E}{\delta v_s(\mathbf{r})} = \int d\mathbf{r}' \frac{\delta E}{\delta n(\mathbf{r}')} \frac{\delta n(\mathbf{r}')}{\delta v_s(\mathbf{r})}$$

- The domain of search within the OEP is augmented compared to the DFT domain.
- This can lead to lower energies than DFT results.

Alternative Approach

M. K. Harbola and V. Sahni, Phys. Rev. Lett. **62**, 489 (1989)



using classical electrodynamics

$$E_x[\rho] = \frac{1}{2} \int \int \frac{\rho(\mathbf{r})\rho_x(\mathbf{r},\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}d\mathbf{r}'$$

corresponding electric field

$$\vec{E}_x = \int \frac{\rho_x(\mathbf{r},\mathbf{r}')(\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} d\mathbf{r}'$$

The potential is the work bringing an electron from infinity to \mathbf{r} :

$$W_x(\mathbf{r}) = - \int_{-\infty}^{\mathbf{r}} \vec{E}_x \cdot d\mathbf{l}$$

Conjecture:

$$W_x(\mathbf{r}) = v_x(\mathbf{r})$$

Summary

- a solution to the self interaction problem (for non-periodic systems) was presented, using only derivatives with respect to the density
- analytic, closed form treatment of the Coulomb energy within Kohn-Sham density functional theory
- an explicit expression for calculating the Coulomb potential, avoiding self-interaction effects by construction
- quantum mechanically correct form of the Coulomb energy using the pair density
- fulfills the 2nd Hohenberg-Kohn theorem: no lower than ground state energy
- formalism and code is developed for all non-periodic systems (so far) (need: all occupied orbitals, boundary conditions for Poisson solver)
- further developments: periodic systems

Reference: Gonis, MD, Nicholson, Stocks, Solid State Communications, 2012, accepted

